

# User Manual for the paper titled “Algorithm xxx: A Suite of Compact Finite Difference Schemes”

MANI MEHRA and KULDIP SINGH PATEL, Indian Institute of Technology Delhi,  
India.

---

Our MATLAB routine have one MATLAB function and three folders.

- `test.m`: This MATLAB function computes the error between analytic differentiation and corresponding compact finite difference approximation for a given function.
- `cfdm_periodic`: This folder has the MATLAB files for computing the compact finite difference approximations for periodic boundary condition in one and two dimensions.
- `cfdm_dirichlet`: This folder has the MATLAB files for computing the compact finite difference approximations for dirichlet boundary condition in one and two dimensions.
- `cfdm_Neumann`: This folder has the MATLAB files for computing the compact finite difference approximations for Neumann boundary in one dimension.

@@

The function `test.m` computes the error between analytic differentiation and corresponding compact finite difference approximation for a given function.

>>test

@@

## **Compact finite difference schemes for periodic boundary in one dimension**

In this case,  $N$  is the number of grid points,  $p$  is the order of accuracy desired,  $x_l$  is the left end of the interval and  $x_r$  is the right end of the interval. Differentiation matrix  $D$  is order of  $N$ .  $f(x)$  is a function defined as  $f = (f_1, f_2, \dots, f_N)^T$ .

1.

`compact_first_periodic.m`

The function `compact_first_periodic.m` computes a differentiation matrix for first derivative approximation. The calling command for this function is

---

Author’s address: M. Mehra (mmehra@maths.iitd.ac.in) and K. S. Patel (kuldip@maths.iitd.ac.in), Indian Institute of Technology Delhi, Hauz Khas, New Delhi-110 016  
Permission to make digital/hard copy of all or part of this material without fee for personal or classroom use provided that the copies are not made or distributed for profit or commercial advantage, the ACM copyright/server notice, the title of the publication, and its date appear, and notice is given that copying is by permission of the ACM, Inc. To copy otherwise, to republish, to post on servers, or to redistribute to lists requires prior specific permission and/or a fee.  
© 20YY ACM 0098-3500/20YY/1200-0001 \$5.00

```
>>[D]=compact_first_periodic(N,p,x_l,x_r)
```

Permissible values for  $p$  are 4, 6, 8 and 10. Then first derivative approximation ( $f'$ ) for any function  $f$  can be written as

$$f' \approx D * f.$$

2.

`compact_second_periodic.m`

The function `compact_second_periodic.m` computes a differentiation matrix for second derivative approximation. The calling command for this function is

```
>>[D]=compact_second_periodic(N,p,x_l,x_r)
```

Permissible values for  $p$  are 4, 6, 8 and 10. Second derivative approximation ( $f''$ ) for any function  $f$  can be written as

$$f'' \approx D * f.$$

3.

`compact_third_periodic.m`

The function `compact_third_periodic.m` computes a differentiation matrix for third derivative approximation. The calling command for this function is

```
>>[D]=compact_third_periodic(N,p,x_l,x_r)
```

Permissible values for  $p$  is 6. Third derivative approximation ( $f'''$ ) for any function  $f$  can be written as

$$f''' \approx D * f.$$

4.

`compact_fourth_periodic.m`

The function `compact_fourth_periodic.m` computes a differentiation matrix for fourth derivative approximation. The calling command for this function is

```
>>[D]=compact_fourth_periodic(N,p,x_l,x_r)
```

Permissible values for  $p$  are 4 and 6. Fourth derivative approximation ( $f''''$ ) for any function  $f$  can be written as

$$f'''' \approx D * f.$$

@@

### Compact finite difference schemes for periodic boundary in two dimensions

In this case,  $N$  is the number of grid points,  $p$  is the order of accuracy desired,  $x_l$  and  $x_r$  are the left end and the right end of the interval in  $x$  direction,  $y_l$  and  $y_r$  are the left end and right end of the interval in  $y$  direction. Differentiation matrix  $D$  is order of  $N^2$  and permissible values for  $p$  are 4, 6, 8 and 10. If  $f(x, y)$  is a function defined as  $f = (f_{1,1}, f_{2,1}, \dots, f_{N,1}, f_{1,2}, f_{2,2}, \dots, f_{N,2}, \dots, f_{1,N}, f_{2,N}, \dots, f_{N,N})^T$ .

5.

`compact_first_periodic_2dx.m`

The function `compact_first_periodic_2dx.m` computes a differentiation matrix for first order partial derivative ( $\frac{\partial}{\partial x}$ ) approximation. The calling command for this function is

```
>>[D]=compact_first_periodic_2dx(N,p,x_l,x_r)
```

First order partial derivative approximation ( $\frac{\partial f}{\partial x}$ ) for any function  $f$  can be written as

$$\frac{\partial f}{\partial x} \approx D * f.$$

6.

`compact_first_periodic_2dy.m`

The function `compact_first_periodic_2dy.m` computes a differentiation matrix for first order partial derivative approximation with respect to  $y$  variable ( $\frac{\partial f}{\partial y}$ ). The calling command for this function is

```
>>[D]=compact_first_periodic_2dy(N,p,y_l,y_r)
```

First order partial derivative approximation ( $\frac{\partial f}{\partial y}$ ) for any function  $f$  can be written as

$$\left(\frac{\partial f}{\partial y}\right) \approx D * f.$$

7.

`compact_second_periodic_2dxx.m`

The function `compact_second_periodic_2dxx.m` computes a differentiation matrix for second order partial derivative approximation. The calling command for this function is

```
>>[D]=compact_second_periodic_2dxx(N,p,x_l,x_r)
```

Second order partial derivative approximation ( $\frac{\partial^2 f}{\partial x^2}$ ) for any function  $f$  can be written as

$$\left(\frac{\partial^2 f}{\partial x^2}\right) \approx D * f.$$

8.

`compact_second_periodic_2dyy.m`

The function `compact_second_periodic_2dyy.m` computes a differentiation matrix for second order partial derivative approximation. The calling command for this function is

```
>>[D]=compact_second_periodic_2dyy(N,p,y_l,y_r)
```

Second order partial derivative approximation  $\left(\frac{\partial^2 f}{\partial y^2}\right)$  for any function  $f$  can be written as

$$\left(\frac{\partial^2 f}{\partial y^2}\right) \approx D * f.$$

9.

`compact_mixed_periodic_2dxy.m`

The function `compact_mixed_periodic_2dxy.m` computes a differentiation matrix for mixed derivative approximation. The calling command for this function is

```
>>[D]=compact_mixed_periodic_2dxy(N,p,x_l,x_r,y_l,y_r)
```

Mixed derivative approximation  $\left(\frac{\partial^2 f}{\partial x \partial y}\right)$  for any function  $f$  can be written as

$$\left(\frac{\partial^2 f}{\partial x \partial y}\right) \approx D * f.$$

@@

#### Compact finite difference schemes for Dirichlet boundary in one dimension

In this case,  $N$  is the number of grid points,  $p$  is the order of accuracy desired,  $x_l$  is the left end of the interval and  $x_r$  is the right end of the interval. Differentiation matrix  $D$  is of order  $N$  and permissible values for  $p$  are 4 and 6.  $f(x)$  is a function defined as  $f = (f_1, f_2, \dots, f_N)^T$ .

10.

`first_compact_dirichlet.m`

The function `first_compact_dirichlet.m` computes a differentiation matrix for first derivative approximation in case of Dirichlet boundary conditions. The calling command for this function is

```
>>[D]=first_compact_dirichlet(N,p,x_l,x_r)
```

First derivative approximation ( $f'$ ) for any function  $f$  can be written as

$$f' \approx D * f.$$

11.

`second_compact_dirichlet.m`

The function `second_compact_dirichlet.m` computes a differentiation matrix for second derivative approximation in case of Dirichlet boundary conditions. The calling command for this function is

```
>>[D]=second_compact_dirichlet(N,p,x_l,x_r)
```

Second derivative approximation ( $f''$ ) for any function  $f$  can be written as

$$f'' \approx D * f.$$

@@

#### Compact finite difference schemes for Dirichlet boundary in two dimensions

In this case,  $N$  is the number of grid points,  $p$  is the order of accuracy desired,  $x_l$  and  $x_r$  are the left end and the right end of the interval in  $x$  direction,  $y_l$  and  $y_r$  are the left end and right end of the interval in  $y$  direction. In this case, differentiation matrix  $D$  is order of  $N^2$ , permissible values for  $p$  are 4 and 6 and  $f(x, y)$  is a function in two variable defined as  $f = (f_{1,1}, f_{2,1}, \dots, f_{N,1}, f_{1,2}, f_{2,2}, \dots, f_{N,2}, \dots, f_{1,N}, f_{2,N}, \dots, f_{N,N})^T$ .

12.

`first_compact_dirichlet_2dx.m`

The function `first_compact_dirichlet_2dx.m` computes a differentiation matrix for first order partial derivative  $(\frac{\partial}{\partial x})$  approximation. The calling command for this function is

```
>>[D]=first_compact_dirichlet_2dx(N,p,x_l,x_r)
```

First order partial derivative approximation  $(\frac{\partial f}{\partial x})$  for any function  $f$  can be written as

$$\frac{\partial f}{\partial x} \approx D * f.$$

13.

`first_compact_dirichlet_2dy.m`

The function `first_compact_dirichlet_2dy.m` computes a differentiation matrix for first order partial derivative approximation with respect to  $y$  variable  $(\frac{\partial f}{\partial y})$ . The calling command for this function is

```
>>[D]=first_compact_dirichlet_2dy(N,p,y_l,y_r)
```

First order partial derivative approximation  $(\frac{\partial f}{\partial y})$  for any function  $f$  can be written as

$$\left(\frac{\partial f}{\partial y}\right) \approx D * f.$$

14.

`second_compact_dirichlet_2dxx.m`

The function `second_compact_dirichlet_2dxx.m` computes a differentiation matrix for second order partial derivative approximation. The calling command for this function is

```
>>[D]=second_compact_dirichlet_2dxx(N,p,x_l,x_r)
```

Second order partial derivative approximation  $(\frac{\partial^2 f}{\partial x^2})$  for any function  $f$  can be written as

$$\left(\frac{\partial^2 f}{\partial x^2}\right) \approx D * f.$$

15.

`second_compact_dirichlet_2dyy.m`

The function `compact_second_periodic_2dyy.m` computes a differentiation matrix for second order partial derivative approximation. The calling command for this function is

```
>>[D]=second_compact_dirichlet_2dyy(N,p,y_l,y_r)
```

Second order partial derivative approximation  $\left(\frac{\partial^2 f}{\partial y^2}\right)$  for any function  $f$  can be written as

$$\left(\frac{\partial^2 f}{\partial y^2}\right) \approx D * f.$$

16.

```
mixed_compact_dirichlet_2dxy.m
```

The function `mixed_compact_dirichlet_2dxy.m` computes a differentiation matrix for mixed derivative approximation in case of Dirichlet boundary conditions. The calling command for this function is

```
>>[D]=mixed_compact_dirichlet_2dxy(N,p,x_l,x_r,y_l,y_r)
```

Mixed derivative approximation for any function  $f$  can be written as

$$\frac{\partial^2 f}{\partial x \partial y} \approx D * f.$$

```
@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@
```

**Compact finite difference schemes for Neumann boundary in one dimension**

17.

```
compact_second_neumann.m
```

The function `compact_second_neumann.m` computes a differentiation matrix for second derivative approximation in case of Neumann boundary conditions. The calling command for this function is

```
>>[D,K]=compact_second_neumann(N,u_left,u_right,x_l,x_r)
```

where  $D$  is of order  $N$ ,  $K$  is a  $N \times 1$  vector, `u_left` is the left Neumann boundary condition and `u_right` is the right Neumann boundary condition. Second derivative approximation ( $f''$ ) for any function  $f$  can be written as

$$f'' \approx D * f + K.$$